
Antibiotic Resistance, Virulence Factors and Phylogenetic Analysis of Efflux Proteins of Coagulase Negative *Staphylococcus* Isolates from Sewage Samples

Kannan Balachander, Deborah Gnana Selvam Alexander *

Post-Graduate Department of Microbiology, The American College, Madurai, India

Email address:

deborahalexander12@gmail.com (D. G. S. Alexander)

*Corresponding author

To cite this article:

Kannan Balachander, Deborah Gnana Selvam Alexander. Antibiotic Resistance, Virulence Factors and Phylogenetic Analysis of Efflux Proteins of Coagulase Negative *Staphylococcus* Isolates from Sewage Samples. *Pharmaceutical Science and Technology*.

Vol. 5, No. 1, 2021, pp. 14-23. doi: 10.11648/j.pst.20210501.13

Received: February 24, 2021; **Accepted:** March 26, 2021; **Published:** May 26, 2021

Abstract: Multidrug-resistant pathogens are now emergent worldwide and pose a serious threat to disease prevention and treatment. The World Health Organization declared anti-microbial resistance as a major peril to human beings. *Staphylococcus* is part of the normal microbiome of humans and are responsible for opportunistic infections. Like *S. aureus*, Coagulase-negative staphylococcus are also clinically important as the causal agents of severe diseases, nosocomial infections, catheter-associated infection, bacteremia, septicemia. Methicillin resistant *Staphylococci* species are widely known and methicillin resistant CoNS, known as MR-CoNS, have been found to carry the *mecA* gene. Various virulence factors help these organisms in their pathogenicity and antibiotic-resistance is aided by the presence of efflux pumps. In our study, we have isolated 45 isolates from sewage water which were tentatively identified as CoNS based on biochemical characteristics. Then virulence and antibiotics susceptibility patterns were screened by standard protocols. Multiple drug resistant CoNS were found in our study and *in silico* analysis of an annotated protein sequence of the efflux pump SepA of *Pseudomonas putida* was carried out. Sequence analysis of the efflux protein gene revealed that it is phylogenetically related to the AcrA of *Staphylococcus* sp. and a RND transporter of *Vibrio* sp. The widespread presence of MR-CoNS is a cause for serious concern as sewage treatment plants are reservoirs for the spread of antibiotic resistance.

Keywords: Coagulase Negative *Staphylococcus*, Efflux Proteins, Multiple Drug Resistant Organism, Phylogenetic Analysis

1. Introduction

The staphylococci are gram-positive microorganisms, the size around 0.5µm- 1 µm in diameter. They are observed as clusters under a microscope. Occasionally it can be seen in short chains. They are ubiquitous, mostly found in human skin and mucus membranes. Staphylococci are salt-tolerant and they can grow in 7.5% NaCl. Staphylococci are often hemolytic. Coagulase test helps to distinguish the *S. aureus* from other staphylococci organisms and some of *S. aureus* from environmental isolates are incapable of producing coagulase. In the genus *Staphylococcus*, the *S. aureus* and *S. intermedius* are the only coagulase-positive organisms and all other species are coagulase-negative. *Staphylococcus aureus* is a common microflora found on our skin surfaces and nasal

passages of healthy human beings. Approximately 25-40% of the human population is colonized with *S. aureus* organism [1]. MR-CoNS harboring the *mecA* gene, which helps resist methicillin and other β- lactam antimicrobials, are referred to as methicillin-resistant *Staphylococcus* (MR-CoNS). Methicillin-resistant *Staphylococcus* (MR-CoNS) is a dangerous pathogen that causes infection in various parts of the human body. It causes mild infections on the skin, like sores, boil, or abscesses, but it can also cause severe infections such as infection of the bloodstream, surgical wounds, the lungs, or urinary tract infections. Most MR-CoNS infections are not serious, some can be life-threatening [2]. Since MR-CoNS infections are difficult to treat. Methicillin-resistant *Staphylococcus aureus* also causes severe infection, which is different from MR-CoNS. *Staphylococcus* resistance to

methicillin was reported soon after its introduction in October 1960 [3]. MR- CoNS is now endemic in India. The incidence of methicillin-resistant *Staphylococcus* varies from 50% in South India [4] to 25% in the western part of India [5]. CA-MRSA has been increasingly reported in India [6]. In India, bloodstream infections (BSI) are caused by *Staphylococcus* species and CoNS are also involved in causing BSI. Those infections are acquired in healthcare settings. *S. epidermidis* are most frequently involved in BSI in developing countries [7].

Antibiotics are widely used to treat numerous infectious diseases. As a result, antibiotic resistant bacteria are also rampant. According to the Centre for Disease Control, at least 2.8 million people in the United States become infected with antibiotic-resistant bacteria each year, with the mortality reaching 35,000 as a direct result of such infections [8]. The role of wastewater as an environmental reservoir in the creation and spread of antibiotic resistance is vital [9]. Particularly when a huge percentage of antibiotics end up in the environment [10]. This includes β -lactam antibiotics, which have been isolated from both soil and drainage systems [11]. In both hospital and wastewater drainage systems, *mecA* gene has been detected [12]. Both culture-based techniques and molecular analysis has found MRSA has been found in municipal water [13]. In the present study, we isolated and identified the CoNS from a municipal Waste Water Treatment Plant WWTP, Madurai, and screened the isolates for multiple antibiotic resistance and compared it with other studies from the locality and other Indian strains. These isolates were tested for antibiotic susceptibility. With the help of sequences obtained from BacMet, an *in silico* sequence analysis of the SepA superfamily of efflux protein was carried out.

2. Material and Methods

2.1. Sample Collection and Processing

The wastewater treatment plant at Avaniyapuram, Madurai was first started in 1924. It receives wastewater from the south zone of the river Vaigai. It is one of the largest plants in Madurai. At present, the plant receives 15 to 20Mld (million liters per day) of wastewater from the main pumping station at Santhaipeitai, and the daily discharge depends upon the availability of electricity and capacity of the machinery. One hundred milliliters of wastewater were collected in autoclaved glass conical flask at Avaniyapuram wastewater plant and the samples were transported within 1h to the laboratory in an icebox. We collected the wastewater sample once in September 2019, December 2019, and February 2020.

2.2. Sample Processing and Isolation

Collected wastewater samples were centrifuged in sterile centrifuge tubes at 8000rpm for 30 minutes at 4°C. Supernatant was discarded and the pellet was collected. The pellet was inoculated into the Nutrient broth with 7.5% NaCl for enrichment process [14] and incubated at 37°C for

overnight incubation. A loopful of culture from the nutrient broth was streaked onto sterile Mannitol Salt Agar and Blood agar plates aseptically and incubated at 37°C for 18-24 h.

2.3. Biochemical Characterization and Detection of Virulence Factors of *Staphylococcus sp.*

The cultured isolates were performed various biochemical tests such as gram staining, coagulase, catalase, oxidase, and TSI test. The virulence factors were screened by various tests such as DNase, hemolysis, starch hydrolysis, and biofilm identification by congo red test.

2.4. Antibiotic Susceptibility Test

Antibiotic susceptibility was performed by the Kirby Bauer method which is a standard diffusion procedure and helps to determine the susceptibility of isolated bacteria to different antibiotics. The filter paper discs are impregnated with antibiotics of specific concentrations. Mueller-Hinton agar plates were used in this procedure and overnight broth cultures of the bacterial isolates were swabbed into the surface of the medium and antibiotic discs were placed on the seeded surface. After incubation, the zone of inhibition was formed. The measuring of the zone of inhibition helps to interpret the results based on the standard chart developed by The Clinical & Laboratory Standards Institute. Twenty-five of the isolates were subjected to antibiotic susceptibility screening.

2.5. Phylogenetic Analysis

SepA is an efflux pump which belongs to the Resistance Nodulation cell Division (RND) superfamily of efflux proteins. It is present in the chromosome of *Pseudomonas sp.* involved in the efflux of multiple drugs. It contains 382 amino acids. This sequence was retrieved from BacMet (antibacterial biocide and metal resistance gene database) and was aligned with other sequences by protein BLAST.

3. Results

The sewage wastewater sample was collected at the Avaniyapuram sewage water treatment plant in a sterile autoclaved conical flask. Before sample collection, pH and temperature were measured *in situ*.

Table 1. pH and temperature of the sample collection from the sewage plant.

Sample collection period	pH	Temperature
September -2019	8.5	37°C
December -2019	9	38.5°C
February -2020	8.5	35.8°C

After overnight incubation, nutrient broth showing turbidity was taken as positive for presumptive staphylococci. Then a loopful of nutrient broth culture was taken and streaked (quadrant streaking) onto the MSA plate and blood agar plate aseptically. Both plates were incubated for 24h at room temperature. After incubation, it was observed that there were too many colonies on the MSA

plate, which turned from red to yellow color showing that the isolates utilize mannitol as a source of carbon. In the blood agar plate, the colonies have grown appeared white and some of the colonies were β- hemolytic. Then 45 single colonies were chosen for maintenance as pure cultures and further characterization. The selected isolates were streaked using sterile inoculation needle onto Nutrient Agar slants supplemented with 7.5% NaCl and after growth was observed, stored at RT.



Figure 1. MSA and Blood agar plate showing the growth of presumptive *Staphylococci*.

Most of the isolates are purple indicating gram-positive organisms. They were found to be cocci and were seen in clusters. Out of the 45 isolates, 41 were found to be gram-positive cocci. $41/45 \times 100 = 91.11\%$. Therefore, 91% of the isolates were gram-positive, cocci, and cluster-forming bacteria and tentatively identified as *Staphylococcus* sp.




Figure 2. Gram staining images under light microscope 100x.

3.1. Biochemical Tests

The preliminary biochemical tests performed were oxidase

and catalase tests. Oxidase test was carried out by disc method which is quick with the results was observed within few seconds. 36 of the isolates are oxidase negative, whereas the other 9 isolates are oxidase-positive which produced purple color after they were smeared on the disc. In catalase test, all of the isolates are produces bubbles, so all isolates are presence of catalase. It is a distinguishing feature that separates *Staphylococci* from *Streptococci* which are catalase negative.



Figure 3. Oxidase, Catalase and TSI test result.

In the tube coagulase test, after inoculation, the tube was observed respectively in 2h, 4h, 6h, 8h, and up to 24h. However, no clotting was observed; all isolates are coagulase-negative. After incubation of TSI agar, color changes were observed in the butt, slant and gas production were observed; of the 25 isolates, 22 isolates utilized 3 sugars and others are not.

3.2. Virulence Factor Test

In the Starch hydrolysis test, following incubation, iodine solution was poured on the surface of the starch medium; zone of clearance representing starch hydrolysis was observed around 2 isolates which represents positive result and all other 43 isolates are negative for amylase. For the detection of DNase production, following incubation, 1N HCl solution was flooded on the surface of DNase medium and observe for the presence of a clear zone around the isolates; all isolates were negative for the presence of DNase. For hemolysis test, after 24 h incubation, the plates were checked for the presence of clear zones; all isolates were negative for hemolysis. Similarly, none of the isolates was capable of biofilm formation which was checked by Congo red plate method.




Figure 4. Starch hydrolysis and Hemolysis activity.

Table 2. Biochemical test results of isolated CoNS organisms.

Colony no	Gram staining	Oxidase	Catalase	Coagulase	DNase	TSI	Starch hydrolysis	Biofilm formation	Hemolysis
1.	Positive	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
2.	Positive	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
3.	Negative	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
4.	Positive	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
5.	Positive	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
6.	Positive	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
7.	Negative	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
8.	Negative	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
9.	Positive	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
10.	Positive	Positive	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
11.	Positive	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
12.	Positive	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
13.	Positive	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
14.	Positive	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
15.	Positive	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
16.	Positive	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
17.	Positive	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
18.	Positive	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
19.	Positive	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
20.	Negative	Negative	Positive	Negative	Negative	-	Negative	Negative	γ - Hemolysis
21.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
22.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
23.	Positive	Positive	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
24.	Positive	Positive	Positive	Negative	Negative	Negative	Positive	Negative	γ - Hemolysis
25.	Positive	Positive	Positive	Negative	Negative	Negative	Negative	Negative	γ - Hemolysis
26.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
27.	Positive	Positive	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
28.	Positive	Positive	Positive	Negative	Negative	Negative	Negative	Negative	γ - Hemolysis
29.	Positive	Positive	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
30.	Positive	Positive	Positive	Negative	Negative	Positive	Positive	Negative	γ - Hemolysis
31.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
32.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
33.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
34.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
35.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
36.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
37.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
38.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
39.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
40.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
41.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
42.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
43.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
44.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis
45.	Positive	Negative	Positive	Negative	Negative	Positive	Negative	Negative	γ - Hemolysis

3.3. Antibiotic Susceptibility Test

Twenty isolates were subjected to antibiotic susceptibility test (Kirby Bauer method). After incubation, the zone inhibition was measured and identified whether they were sensitive or resistant based on CLSI standards. All 20 isolates are resistant to methicillin; 6 isolates are resistant to

norfloxacin and the other 14 were sensitive. One culture was resistant to chloramphenicol and the other 19 are sensitive. Thirteen isolates were sensitive to ciprofloxacin, 6 are intermediate and one was resistant. All isolates were sensitive to tetracycline and levofloxacin.




Figure 5. Antibiotic Susceptibility Test (Kirby Bauer method).

Table 3. Antibiotic Resistance Profile of CoNS isolated from sewage water samples.

Colony no	Methicillin (5µg)	Tetracycline (30 µg)	Norfloxacin (10 µg)	Levofloxacin (5 µg)	Chloramphenicol (30 µg)	Ciprofloxacin (5 µg)
1.	R	S	S	S	S	S
2.	R	S	S	S	S	S
3.	R	S	S	S	S	S
4.	R	S	S	S	S	S
5.	R	S	S	S	S	S
6.	R	S	R	S	R	I
7.	R	S	S	S	S	S
8.	R	S	S	S	S	I
9.	R	S	S	S	S	I
10.	R	S	S	S	S	S
11.	R	S	S	S	S	S
12.	R	S	R	S	S	I
13.	R	S	R	S	S	R
14.	R	S	R	S	S	I
15.	R	S	R	S	S	I
16.	R	S	S	S	S	S
17.	R	S	S	S	S	S
18.	R	S	R	S	S	S
19.	R	S	R	S	S	S
20.	R	S	S	S	S	S

R- Resistant, S- Sensitive, I- Intermediate.

3.4. Phylogenetic Analysis

Sequence analysis of the SepA (of *Pseudomonas putida*) efflux protein with other sequences revealed that it is homologous to various efflux proteins like AcrA of *Staphylococcus sp.* The tree shows that this SepA protein is

also homologous to another resistance efflux protein called RND transporter of *Vibrio sp.* The phylogenetic analysis showed that these proteins have evolved from a common ancestral gene.



Figure 6. Phylogenetic tree showing the relationship between *SepA* probable efflux pump periplasmic linker protein of *Pseudomonas putida* with the multidrug-efflux RND transporter periplasmic adaptor subunit *AcrA* of *Staphylococcus* sp.

4. Discussion

The CoNS are mostly associated with clinical device infections as they readily form biofilms on the surface of invasive devices like a catheter. The CoNS are important in nosocomial infections and other clinical infections. CoNS are ubiquitous; they are mostly found on skin surfaces and mucus membranes. Most of them are opportunistic pathogens. These organisms can be found as normal microflora on every human being. Staphylococci are often hemolytic and cause disease in humans and animals. Mostly they cause wound infections, bloodstream infections, and hospital-acquired infections such as surgical wound infections. The CoNS *S. epidermidis* causes the infection associated with indwelling medical devices. The *S. saprophyticus* causes UTI infections in girls. The other Staphylococcal organisms such as *S. warneri*, *S. haemolyticus*, *S. intermedius*, and *S. schleiferi* are also pathogenic which cause diseases infrequently in humans [15, 16].

Sewage water includes a wide variety of microorganisms such as bacteria, viruses, and protozoa. Most microorganisms in wastewater are pathogenic and highly virulent. Wastewater can contain a lot of opportunistic pathogens (e.g., *Escherichia coli*, *Enterobacter cloacae*, *Enterococcus faecalis*, *Klebsiella pneumoniae*, *Proteus vulgaris*, or *Pseudomonas aeruginosa*) which can cause various systemic infections, particularly among people with poor and impaired immune systems. *Salmonella* and *Shigella*, or enteropathogenic strains of *Escherichia coli*, responsible for gastroenteritis, salmonellosis, shigellosis, can also be present in wastewater [16, 17].

CoNS *S. pasteurii* was identified in a significant proportion (65.7%) of drinking water supply *S. carnosus*, *S. condimenti*, *S. equorum*, *S. piscifermentans*, *S. succinus*, and *S. xylosus* representing staphylococcal organisms linked to fermented foods [18]. *S. epidermidis* and other CoNS are generally known as non-pathogenic having little virulence factors. The most frequently appearing infectious agent is the incremental clarification of microbial pathogenesis related to the involvement of foreign species in the organism and especially concerning *S. epidermidis* [19]. Staphylococci are the leading cause of hospital and community-acquired infections, particularly antibiotic-resistant strains. These bacteria are ubiquitous and commonly spread in a wide range of environments, with exceptional survival. In addition to serving as a source for the diffusion of resistance genes to various bacteria, including pathogens, they can survive for long periods in diverse environments. Their capacity for survival and gain resiliency genes can partially enlighten why they remain to be a noteworthy human and animal pathogen [2].

CoNS can be characterized by various tests, such as colony morphology, catalase test, gram staining, slide and tube coagulase test (i.e., bound coagulase and free coagulase) (read after 4, 24 h) and mannitol salt agar, deoxyribonuclease

test. The biochemical tests include novobiocin sensitivity, mannitol utilization, ornithine decarboxylation, urease production, phosphatase test, polymyxin B disc test, Voges–Proskauer test, mannose fermentation, trehalose fermentation [20]. In our study most of the above-mentioned tests were performed for the tentative identification of CoNS.

Bacteria produce extracellular enzymes or exoenzymes, which help them in the invasion of the host. These enzymes act as virulence factors making the strains pathogenic. During an infection, bacterial survival is a mechanism dependent on the organism's ability to evade and bypass host defense strategies. *S. aureus* synthesizes a large number of toxins and exoproteins. Nearly all synthesizes a lot of enzymes and cytotoxins, including four hemolysins (alpha, beta, gamma, and delta), collagenase, lipases, nucleases, hyaluronidase, and proteases [21]. These proteins can principally have the role of transforming host tissues into growth nutrients necessary for the survival of the invading bacteria. Some strains also synthesize other extra exoproteins such as the toxic shock syndrome toxin-1 (TSST 1) (SEA, SEB, SEC, SED, SEE, SEG, SEH, and SEI), and the toxins A and B. Gamma-hemolysine is involved in the lysis of mammalian erythrocytes. These toxins also affect neutrophils and macrophages. Coagulase produced by *Staphylococcus* sp. Interferes with blood clotting by binding prothrombin to a complex called staphylothrombin in the host. These are some of the strategies adapted by *S. aureus* cells defend themselves from the host's immune response [22].

Hydrolytic amylases (α -amylase, β -amylase, and glucoamylase) of starch are among the most widely used in modern biotechnology. Fungi and bacteria secrete amylases to convert excess extracellular starch into sugars. *Staphylococcus aureus* is well known as a major human Gram-positive pathogen that produces multiple surface and secretory proteins, including separate enzymes and pathogenic factors, supporting colonization and host-tissue infection. α -amylase is an enzyme separated from *S. aureus* that catalyzes the colonization and survival [23].

Hemolysin is one of the virulence factors observed in both coagulase-positive and coagulase-negative staphylococci (CoNS). There were substantial differences in non-hemolytic isolates between CoNS, CoPS, and all non-hemolytic isolates are part of CoNS. Most of the *Staphylococcus* isolated from cattle exhibited delta hemolysin on sheep's blood agar plate, while most humans developed alpha-hemolysin. 10 (50%) of the 20 human isolates developed double hemolysin (DH), while the DH-ratio for bovine isolates (one out of 20) was slightly low [24].

In our studies, we screened all CoNS isolates for hemolytic activity in the blood agar supplemented with human blood. All isolates weren't able to produce hemolysis; so, all are non-hemolytic CoNS.

In recent decades, CoNS has been continuously decreasing resistance to most of the antibiotics accessible. In specific penicillin, oxacillin, ciprofloxacin, clindamycin, erythromycin, or gentamicin have established significant and

radical changes in the quantity of resistant isolates [25]. In the 1980s publications comparing *S. epidermis* and *S. haemolyticus* found increase in the number of methicillin-resistant isolates of both species; although MR *S. haemolyticus* strains had a higher MIC [26]. A clear increase in MR-CoNS percentage was witnessed. The steep increase in CoNS infections was observed in a twenty year study in Switzerland as the rate of MR-CoNS isolates recovered from burn patients raised from 11% to 50 percent from 1986 to 2005 [27]. In the United States and four European countries, the proportion of CoNS resistant to oxacillin ranged from 51.4% in France to 75.2% in the United States while 2,905 CoNS isolates were examined in 2001 from SSTIs of hospitalized patients [28].

The *mec* genes are seen in the chromosome by a mobile SCC_{mec} genetic feature. This cassette consists of three main elements: the *mec* gene complex, the *ccr* gene complex, and the joining area ('junkyard' or J). The *mec* gene complex comprises of the *mecA* gene itself, and the *mecI* (repressor), *mecR1* (sensor inducer), and IS431*mec* insert sequence [29]. [30].

In our studies, we performed the antibiotic susceptibility test by disc diffusion method (Kirby-Bauer) for 20 isolates. All 20 isolates are resistant to methicillin (5µg) antibiotic. The 2 isolates are resistant to more than 2 antibiotics. Isolate number 6 resist methicillin, norfloxacin (10 µg), and chloramphenicol (30 µg). The isolate number 13 resists methicillin, norfloxacin (10 µg), and ciprofloxacin (5 µg). Multi-resistant CoNS isolates, displaying simultaneously resistance against at least three antibiotic classes, were prevalent.

The CoNS *S. epidermidis* strain was resistant to 6 distinct antibiotics (AMP, CRO, K, OB, OFX, and OT) [31].

In another study, 16 (29%) staphylococci isolates were resistant to 3 or more antibiotics [32]. In 25 percent of the staphylococcal isolates from patients with chronic blepharitis, some others observed resistance to two or more antibiotics. Multiple antibiotic resistance can be seen as a reaction to extended therapy. Therefore, it is a cause for alarm to detect one strain immune to all antibiotics tested. These strains are a formidable challenge to hospitalized immune-compromised patients. Multidrug resistant CoNS often colonizes the skin of hospital personnel thus exposing vulnerable patients to life threatening infections [33].

Efflux pumping inhibitors (EPIs) may be a powerful strategy to overwhelm MDR. 1-(1-naphthyl-methyl)-piperazine (NMP) and phenylalanine-arginine Î2 naphthylamide (PAÎ2N) are model EPIs that can block the activity AcrB, a large *Escherichia coli* efflux pump, or related homologous resistance pumps -nodulation-cell class division [34].

In *Staphylococcus* spp. efflux pumps is an important resistance mechanism. In addition to the chromosome pump and the plasmid-encoded NorA pump, which has been described previously, *S. aureus* chromosomal pumps include AbcA, NorB, NorC, MepA, MdeA, SdrM and Tet, SmR, and Sav-1866 type MFS [35, 36].

In the present study, the phylogenetic tree proves the SepA

of *P. putida* is homologous with the various efflux proteins seen in *Staphylococcus* sp. And *Vibrio* sp.

5. Conclusion

Currently, antimicrobial resistance microorganisms have increased rampantly worldwide, which is a serious threat to humans and other animals. World Health Organization declared anti-microbial resistance as a catastrophe. These antimicrobial resistance pathogens are frequently found in different environments and often related to anthropogenic activities. Sewage treatment plants are treating and recycling the used contaminated waters. This recycled water is used for again human use. If those multiple drug resistant organisms are present after treatment, this will cause severe threat to human beings. So, we need to develop a modern treatment procedure for those sewage treatment plants that will effectively reduce multiple drug pathogens and organisms. The RND efflux proteins help to resist the antibiotics; they are superfamily efflux proteins involved in multiple drug-resistance. The widespread presence of methicillin resistance in CoNS is a cause for grave concern as sewage treatment plants are reservoirs for the spread of antibiotic resistance. Our study has shed light on the extensive presence of methicillin- CoNS in local sewage treatment plant.

Conflict

The authors declare that they have no competing interests.

Funding

Self-funded project.

Acknowledgements

Authors acknowledge the Post Graduate Department of Microbiology, The American College, Madurai for laboratory facilities and encouragement.

References

- [1] Mangi, Shahida & Hussain, Anwar & Qazi, Muneer & Kanhar, Nisar. (2016). Incidence of multidrug-resistant coagulase-negative Staphylococci in clinical samples of Tertiary care hospitals of Khairpur and Sukkur cities of Pakistan. *International Journal of Biosciences*. 9. 261-269. 10.12692/ijb/9.4.261-269.
- [2] Abulreesh, H. H. (2011). Multidrug-resistant staphylococci in the environment. In *International Conference on Biotechnology and Environment Management* (Vol. 18, pp. 1-6).
- [3] Jevons, M. P., & Parker, M. T. (1964). The evolution of new hospital strains of *Staphylococcus aureus*. *Journal of clinical pathology*, 17(3), 243-250. <https://doi.org/10.1136/jcp.17.3.243>.

- [4] Gopalakrishnan, R., & Sureshkumar, D. (2010). Changing trends in antimicrobial susceptibility and hospital-acquired infections over an 8 years period in a tertiary care hospital in relation to introduction of an infection control programme. *The Journal of the Association of Physicians of India*, 58 Suppl, 25–31.
- [5] Patel, A. K., Patel, K. K., Patel, K. R., Shah, S., & Dileep, P. (2010). Time trends in the epidemiology of microbial infections at a tertiary care center in west India over last 5 years. *J Assoc Physicians India*, 58 (58), 37-40.
- [6] D'Souza, N., Rodrigues, C., & Mehta, A. (2010). Molecular characterization of methicillin-resistant *Staphylococcus aureus* with emergence of epidemic clones of sequence type (ST) 22 and ST 772 in Mumbai, India. *Journal of clinical microbiology*, 48 (5), 1806–1811. <https://doi.org/10.1128/JCM.01867-09>.
- [7] Rathanin Seng, U. L. (2017). High prevalence of methicillin-resistant coagulase-negative staphylococci isolated from a university environment in Thailand. *International Microbiology*, 20 (2), 65-73. doi: 10.2436/20.1501.01.286.
- [8] CDC, Antibiotic/Antimicrobial resistance report in US (2019), Retrieved from; <https://www.cdc.gov/drugresistance/index.html>.
- [9] Martinez, J. L. (2006). Role of non-clinical environments in the selection of virulence and antibiotic resistance determinants in pathogenic bacteria. *J. Biol. Sci*, 6 (1), 1-8.
- [10] Kümmerer, K., Al-Ahmad, A., & Mersch-Sundermann, V. (2000). Biodegradability of some antibiotics, elimination of the genotoxicity and affection of wastewater bacteria in a simple test. *Chemosphere*, 40 (7), 701-710.
- [11] Watkinson, Andrew & Murby, E. J. & Kolpin, Dana & Costanzo, Simon. (2009). The occurrence of antibiotics in an urban watershed: From wastewater to drinking water. *The Science of the total environment*. 407. 2711-23. 10.1016/j.scitotenv.2008.11.059.
- [12] Volkmann, H., Schwartz, T., Bischoff, P., Kirchen, S. and Obst, U. (2004) Detection of clinically relevant antibiotic resistance genes in municipal wastewater using real-time PCR (Taqman). *J Microbiol Methods* 56, 277–286.
- [13] Börjesson, S., Matussek, A., Melin, S., Löfgren, S., & Lindgren, P. E. (2010). Methicillin-resistant *Staphylococcus aureus* (MRSA) in municipal wastewater: an uncharted threat?. *Journal of applied microbiology*, 108 (4), 1244–1251. <https://doi.org/10.1111/j.1365-2672.2009.04515.x>.
- [14] Cyprowski M, Szarapińska-Kwaszewska J, Dudkiewicz B, Krajewski JA, Szadkowska-Stańczyk I (2005) Exposure assessment to harmful agents in workplaces in sewage plant workers. *Med Pr* 56 (3): 213–222.
- [15] Gerardi MH, Zimmerman MC (2005) *Wastewater pathogens*. Wiley, New Jersey.
- [16] Becker, K., Heilmann, C., & Peters, G. (2014). Coagulase-negative staphylococci. *Clinical microbiology reviews*, 27 (4), 870-926.
- [17] Longauerova, A. (2006). Coagulase negative Staphylococci and their participation in pathogenesis of human infections. *Bratislavské lekárske listy*. 107. 448-52.
- [18] Kondapaneni, Saileela & Vanaparthi, Nagaraju & Anukolu, Ravishankarreddy & Kumar, A. N. (2017). Biochemical Characterization and Antibiotic Susceptibility of Coagulase Negative Staphylococci Isolates from Exudates and Body Fluids. *Indian Journal of Mednodent and Allied Sciences*. 5. 10.5958/2347-6206.2017.00012.7.
- [19] Dinges, M. M.; Orwin, P. M.; Schlievert, P. M. Exotoxins of *Staphylococcus aureus*. *Clin. Microbiol. Rev.* 2000, 13, 16–34.
- [20] Cotar, Ani & Chifiriuc, Mariana & Dinu, Sorin & Bucur, Marcela & Iordache (Curutiu), Carmen & Banu, Otilia & Dracea, Olguta & Larion, Cristina & Lazar, Veronica. (2010). Screening of Molecular Virulence Markers in *Staphylococcus aureus* and *Pseudomonas aeruginosa* Strains Isolated from Clinical Infections. *International journal of molecular sciences*. 11. 5273-91. 10.3390/ijms11125273.
- [21] Lakshmi, H. P., Prasad, U. V., Yeswanth, S., Swarupa, V., Prasad, O. H., Narasu, M. L., & Sarma, P. V. (2013). Molecular characterization of α -amylase from *Staphylococcus aureus*. *Bioinformation*, 9 (6), 281–285. <https://doi.org/10.6026/97320630009281>.
- [22] Moraveji, Z., Tabatabaei, M., Shirzad Aski, H., & Khoshbakht, R. (2014). Characterization of hemolysins of *Staphylococcus* strains isolated from human and bovine, southern Iran. *Iranian journal of veterinary research*, 15 (4), 326–330.
- [23] Kresken M, Hafner D. 1999. Drug resistance among clinical isolates of frequently encountered bacterial species in central Europe during 1975–1995. *Infection* 27 (Suppl 2): S2–S8. doi: 10.1007/BF02561661.
- [24] Hamilton-Miller JM, Iiffe A. 1985. Antimicrobial resistance in coagulase-negative staphylococci. *J. Med. Microbiol.* 19: 217–226. doi: 10.1099/00222615-19-2-217.
- [25] Guggenheim M, Zbinden R, Handschin AE, Gohritz A, Altintas MA, Giovanoli P. 2009. Changes in bacterial isolates from burn wounds and their antibiograms: a 20-year study (1986-2005). *Burns* 35: 553–560. doi: 10.1016/j.burns.2008.09.004.
- [26] Jones ME, Karlowsky JA, Draghi DC, Thornsberry C, Sahn DF, Nathwani D. 2003. Epidemiology and antibiotic susceptibility of bacteria causing skin and soft tissue infections in the U.S.A. and Europe: a guide to appropriate antimicrobial therapy. *Int. J. Antimicrob. Agents* 22: 406–419. doi: 10.1016/S0924-8579(03)00154-7.
- [27] Tesch W, Ryffel C, Strässle A, Kayser FH, Berger-Bächli B. 1990. Evidence of a novel staphylococcal mec-encoded element (mecR) controlling expression of penicillin-binding protein 2'. *Antimicrob. Agents Chemother.* 34: 1703–1706. doi: 10.1128/AAC.34.9.1703.
- [28] Hiramatsu K, Katayama Y, Yuzawa H, Ito T. 2002. Molecular genetics of methicillin-resistant *Staphylococcus aureus*. *Int. J. Med. Microbiol.* 292: 67–74. doi: 10.1078/1438-4221-00192.
- [29] Virdis, S., Scarano, C., Cossu, F., Spanu, V., Spanu, C., & De Santis, E. P. (2010). Antibiotic Resistance in *Staphylococcus aureus* and Coagulase Negative Staphylococci Isolated from Goats with Subclinical Mastitis. *Veterinary medicine international*, 2010, 517060. <https://doi.org/10.4061/2010/517060>.
- [30] Pinna A, Zanetti S, Sotgiu M, et al., Identification and antibiotic susceptibility of coagulase negative staphylococci isolated in corneal/external infections. *British Journal of Ophthalmology* 1999; 83: 771-773.

- [31] Archer GL, Climo MW. Antimicrobial susceptibility of coagulase-negative staphylococci. *Antimicrob Agents Chemother* 1994; 38: 2231–7.
- [32] Soparkar, K., Kinana, A. D., Weeks, J. W., Morrison, K. D., Nikaido, H., & Misra, R. (2015). Reversal of the Drug Binding Pocket Defects of the AcrB Multidrug Efflux Pump Protein of *Escherichia coli*. *Journal of bacteriology*, 197 (20), 3255–3264. <https://doi.org/10.1128/JB.00547-15>.
- [33] Otto, M. (2013). Coagulase-negative staphylococci as reservoirs of genes facilitating MRSA infection. *Bioessays* 35, 4-11.
- [34] Xu Z, M. H. (2015). Antibiotic resistance and *mecA* characterization of coagulase-negative staphylococci isolated from three hotels in London, UK. *Front Microbiol*, 6, 947. doi: 10.3389/fmicb.2015.00947.
- [35] Zhang M, O. M. (2011). Prevalence of antiseptic-resistance genes in *Staphylococcus aureus* and coagulase-negative staphylococci colonising nurses and the general population in Hong Kong. *J Hosp Infect.*, 78 (2), 113-117. doi: 10.1016/j.jhin.2011.02.018.
- [36] Li, X. Z., & Nikaido, H. (2009). Efflux-mediated drug resistance in bacteria: an update. *Drugs*, 69 (12), 1555–1623. <https://doi.org/10.2165/11317030-000000000-00000>.